

Percentage of Children Under-Five Who Had Diarrhea in the Last Two Week Preceding the Survey and Treated with ORT by Type and Selected Background Characteristics

Background Characteristics	Type of Treatment				Number of Children
	Home Made Solution	ORS	Both	Neither	
Age of child					
< 6 months	5.6	27.6	0.0	66.8	68
6-11	10.5	20.4	6.9	62.2	142
12-17	7.1	19.3	5.9	67.7	122
18-23	3.8	36.2	1.4	58.6	84
24-59	3.9	14.7	6.0	75.4	154
Sex					
Male	5.7	21.4	4.3	68.6	309
Female	7.2	22.3	5.4	65.1	261
Place of Residence					
West Bank	6.2	18.4	4.1	71.3	400
Gaza Strip	6.8	29.8	6.5	56.9	170
Total	6.4	21.8	4.8	67.0	570

Percentage of Children 12-23 Months Whom Immunization Card Was Checked and Received Specific Immunization Doses, by Selected Background Characteristics

Background Characteristics	Percent of Children Immunized			
	BCG	DPT3/OPV3	Measles	All Doses*
Sex				
Male	42.1	93.2	47.7	79.6
Female	44.4	96.6	50.1	82.9
Place of Residence				
West Bank	21.6	93.8	28.7	75.4
Gaza Strip	90.5	97.3	93.2	94.0
Total	43.2	94.9	48.9	81.3

* Includes (Measles or MMR) and Three Doses of DPT and OPV.

Contraceptives Prevalence (Modern Methods, Traditional Methods) Among Currently Married Women by Type of Contraceptive

Method	Knowledge	Usage	
	Knows the Method	Ever-used	Currently using
Any Method	99.6	66.9	45.3
<u>Any Modern Method</u>	99.6	53.7	30.7
Pill	98.8	24.9	3.8
IUD	99.2	39.2	21.6
Injections	65.1	2.0	0.6
Condoms	68.1	8.9	1.5
Vaginal Methods*	54.3	5.2	0.5
Other	85.6	3.0	2.8
<u>Any Traditional Method</u>	96.9	34.5	14.5
Safe Period	74.2	16.0	5.1
Withdrawal	72.1	15.0	5.8
Breast feeding	93.0	14.1	3.3

* Vaginal Methods Include: Diaphragm, Sperm Foam, Jelly, Foaming and Cream.

Percent Distribution of Currently Married Women, Who Are Using Modern Contraceptive Methods by Source and Type

Source of Method	Type of Modern Method			Total
	Pill	IUD	Other	
Public Hospital	—	3.6	24.2	6.7
Private Hospital	2.4	5.0	27.2	8.5
Public Health Center	3.0	2.6	0.9	2.4
Private Health Center	3.4	7.8	2.3	6.3
Health Center/Maternal Center	16.2	13.8	8.0	13.1
Family Planing Clinic	5.7	13.9	4.3	11.3
Private Doctor	40.7	53.3	12.2	44.7
Pharmacy	28.6	—	20.9	7.0
Total	100	100	100	100

Percent Distribution of Currently Married Non-Pregnant Women Who Are Not Using Any Contraceptive Method and Who Do Not Intend to Use Any, by Reason for Not Intending to Use

Reasons For Not Intending to Use	Place of Residence		Total
	West Bank	Gaza Strip	
Religious Reasons	6.2	5.9	6.1
Opposes Family Planning	8.9	12.7	10.5
Opposition of Husband	11.5	19.3	14.8
Opposition of Relatives	1.3	0.9	1.1
Fear of Side Effects	8.9	9.1	9.0
Lack of Knowledge	1.6	2.0	1.8
Cost Too Much	0.4	0.6	0.5
In Convenient	3.4	2.7	3.1
Menopause	37.1	35.3	36.3
Don't Know	20.7	11.5	16.8
Total	100	100	100
Number of Women	404	284	688

Percentage of Persons Who Had An Unexpected Illness or Injury During the Last Two Weeks Preceding the Survey by Selected Background Characteristics

Background Characteristics	Percent of Persons Who Had Unexpected Illness	Percent of Persons Who Had An Injury	Total Population
Age (Years)			
0-4	35.2	1.5	4469
5-14	7.1	0.7	7045
15-29	7.6	0.8	6941
30-39	10.7	1.0	2530
40-49	11.6	1.0	1690
50+	11.9	1.8	2554
Sex			
Male	13.2	1.3	12869
Female	13.6	0.7	12360
Place of Residence			
West Bank	13.5	1.2	17297
Gaza Strip	13.1	0.6	7932
Total	13.4	1.0	25229

**Number of Household Members with Disability and Disability Rate Per 100,000
Persons, by Sex and Age**


Age Group	Household Population	No. of Disabled Population	Disability Rate/100,000
Male			
0-4	2275	24	1054.9
5-14	3601	82	2277.1
15-64	6555	169	2578.6
65+	439	33	7517.1
Female			
0-4	2195	37	1685.6
5-14	3444	57	1655.1
15-64	6242	112	1780.9
65+	477	27	5660.4
Total			
0-4	4470	61	1364.7
5-14	7045	139	1973.0
15-64	12797	281	2195.8
65+	916	60	6550.2

Percent Distribution of Individuals by Place of Residence and Type of Health Insurance


Type of Health Insurance	West Bank *		Gaza Strip		Total	
	Number	%	Number	%	Number	%
Government	6498	37.6	3314	41.8	9812	38.9
UNRWA	2223	12.9	2433	30.7	4656	18.5
Social Security	170	1.0	140	1.8	310	1.2
Private	560	3.2	198	2.5	758	3.0
Without Insurance	7843	45.3	1847	23.3	9690	38.4
Total	17297	100	7932	100	25229	100

* 3 Cases with missing value in the West Bank.


Percentage of Children Under-five Who Had Diarrhea and Acute Respiratory Tract Infection During the Two Weeks Preceding the Survey by Place of Residence




Percentage of Children Under Five Who Classified Under Nourished According to Anthropometric Measures of Nutritional Status by Place of Residence and Sex




Mean Duration of Breastfeeding Among Children (Last Two) by Selected Background Characteristics




**Percent of Children (2-11) Months Who Had Received Vitamins A/D Supplements
by Place of Residence and Sex**




Percentage of Pregnant Women During the Five Years Preceding the Survey Who Received Pre-natal Care by Place of Residence




**Percent Distribution of Births (Last Two) During the Five Years Preceding the Survey
by Place of Delivery, Place of Residence and Mother's Education**



Percentage of Currently Pregnant Women Who Had Received Tetanus Toxoid by Doses and Place of Residence



Percent Distribution of Births (Last Two) During the Five Years Preceding the Survey Whose Mothers Had Received Post-natal Care by Personnel Providing Post-natal Care



**Contraceptives Prevalence (Modern or Traditional Methods) Among Currently Married Women
by Mother's Education and Place of Residence**

